

Quasilocal mass and surface Hamiltonian in spacetime

Mu-Tao Wang

*Department of Mathematics, Columbia University,
New York, NY 10027, USA*

**E-mail: mtwang@math.columbia.edu*

We discuss the concepts of energy and mass in relativity. On a finitely extended spatial region, they lead to the notion of quasilocal energy/mass for the boundary 2-surface in spacetime. A new definition was found in²⁷ that satisfies the positivity, rigidity, and asymptotics properties. The definition makes use of the surface Hamiltonian term which arises from Hamilton-Jacobi analysis of the gravitation action. The reference surface Hamiltonian is associated with an isometric embedding of the 2-surface into the Minkowski space. We discuss this new definition of mass as well as the reference surface Hamiltonian. Most of the discussion is based on joint work with PoNing Chen and Shing-Tung Yau.

Keywords: Quasilocal mass, surface Hamiltonian.

1. Energy of matter fields and conservation

Relativity is a unified theory of space and time. The spacetime of special relativity is the Minkowski space $\mathbb{R}^{3,1} = \mathbb{R} \times \mathbb{R}^3$ with Lorentz metric of signature $(-1, 1, 1, 1)$. We normalize the speed of light to be 1. The light cone consists of four vectors (t, x, y, z) with $-t^2 + x^2 + y^2 + z^2 = 0$. As nothing travels faster than light, a material particle or an observer moves in future timelike direction.

To each matter field, an energy-momentum tensor T is attached. T is derived from the Lagrangian of the field and is described by first derivatives of the field. In particular, it is a symmetric $(0, 2)$ tensor $T_{\mu\nu}$ which satisfies the conservation law

$$\nabla^\mu T_{\mu\nu} = 0. \quad (1)$$

Without gravitation, the energy of a physical system Ω is obtained by integrating T on Ω with respect to an observer. To be more precise, given a spacelike bounded region Ω , the energy intercepted by Ω as seen by the observer t^ν is the flux integral

$$\int_{\Omega} T_{\mu\nu} t^\mu u^\nu$$

where u^ν is the future timelike unit normal of Ω . The dominant energy condition guarantees $T_{\mu\nu} t^\mu u^\nu \geq 0$.

Suppose t^μ is a constant future-directed timelike unit vector in $\mathbb{R}^{3,1}$. By conservation law (1), $T_{\mu\nu}t^\mu$ is divergence free and thus is dual to a closed 3-form in $\mathbb{R}^{3,1}$, which in turn is $d\omega$ for a 2-form ω . Therefore, $\int_\Omega T_{\mu\nu}t^\mu u^\nu = \int_{\partial\Omega} \omega$ is a linear expression in t^μ . Minimizing among all such observers t^μ gives the quasilocal mass which depends only on the boundary 2-surface $\Sigma = \partial\Omega$. Moreover, $\int_\Omega T_{\mu\nu}u^\nu$ defines a quasilocal energy-momentum 4-vector. This is the prototype of quasilocal mass and quasilocal energy-momentum.

2. Energy in General relativity

In general relativity, spacetime is a 4-dimensional manifold with a Lorentz metric g , the gravitational field. Local causal structure of spacetime remains the same, and each tangent space is isometric to the Minkowski space. Gravitational force is represented by the spacetime curvature of g . The relation between the gravitation field and matter fields is exactly described by the Einstein equation

$$Ric - \frac{1}{2}Rg = 8\pi T \quad (2)$$

where Ric is the Ricci curvature, and R is the scalar curvature of g , respectively. T represents the energy-momentum tensor of all matter fields. This is the Euler-Lagrange equation of the Hilbert-Einstein action.

Concerning energy, one seeks for an energy momentum tensor for gravitation. However, it turns out first derivatives of g are all coordinate dependent, and thus there is no density for gravitational energy. This is Einstein's equivalence principle. One can still integrate T on the right hand side of (2) but this gives only the energy contribution from matters. Indeed, there exists vacuum spacetime, i.e. $T = 0$, with nonzero energy such as Schwarzschild's or Kerr's solution of Einstein's equation. This is gravitational energy by the sheer presence of spacetime curvature. Even without energy density, one can still ask the question: what is the energy in a system Ω , counting contribution from gravitation and all matter fields?

In special relativity, the energy integral of T on Ω depends only on the boundary data by energy conservation. One expects energy conservation in general relativity as well, and thus this information should be encoded in the geometry the two-dimensional boundary surface $\Sigma = \partial\Omega$.

This leads to the well-known problem of quasilocal energy/mass in general relativity. The first one in Penrose's 1982 list¹⁸ of major unsolved problems in classical general relativity is "Find a suitable quasilocal definition of energy-momentum in general relativity".

3. Total energy and mass

Einstein's field equation is derived from variation of the Einstein-Hilbert action on a spacetime domain M :

$$\frac{1}{16\pi} \int_M R + \frac{1}{8\pi} \int_{\partial M} K + \int_M L(g, \Phi)$$

where K is the trace of the second fundamental form of ∂M and Φ represents all the matter fields.

Formally applying Hamilton-Jacobi analysis to this action, we obtain $T^*_{\mu\nu}$, the so called Einstein pseudo tensor, which is expressed in terms of first derivatives of g and satisfies $\nabla^\mu T^*_{\mu\nu} = 0$.

Here is Hermann Weyl's (1921) comment on $T^*_{\mu\nu}$ (the English translation is quoted from⁸):

"Nevertheless it seems to be physically meaningless to introduce the $T^*_{\mu\nu}$ as energy components of the gravitational field; for, these quantities are neither a tensor nor are they symmetric. In fact by choosing an appropriate coordinate system all the $T^*_{\mu\nu}$ can be made to vanish at any given point; for this purpose one only needs to choose a geodesic (normal) coordinate system. And on the other hand one gets $T^*_{\mu\nu} \neq 0$ in a 'Euclidean' completely gravitationless world when using a curved coordinate system, but where no gravitational energy exists. Although the differential relations ($\nabla^\mu T^*_{\mu\nu} = 0$) are without a physical meaning, nevertheless by integrating them over an *isolated system* one gets invariant conserved quantities".

An isolated system is modeled on an unbounded and asymptotically flat space-time where gravitation is weak at infinity. There are two notions of total mass associated with such a system, one at spatial infinity and the other at null infinity.

Arnold-Deser-Misner¹ applied Hamilton-Jacobi analysis of the Einstein-Hilbert action to such a system that is asymptotically flat at spatial infinity, and obtained a total energy-momentum that is conserved.

Suppose (Ω, g_{ij}, p_{ij}) is asymptotically flat, i.e. there is a compact subset K of Ω such that $\Omega \setminus K$ is a finite union of ball complements in \mathbb{R}^3 , and on each component there is asymptotically flat coordinate system such that $g_{ij} - \delta_{ij} \sim 0$ and $p_{ij} \sim 0$ with appropriate decay rate on their derivatives. The total energy is

$$E = \lim_{r \rightarrow \infty} \frac{1}{16\pi} \int_{S_r} (\partial_j g_{ij} - \partial_i g_{jj}) dv^i,$$

where S_r is the coordinate sphere of coordinate radius r . The total momentum is

$$P_k = \lim_{r \rightarrow \infty} \frac{1}{16\pi} \int_{S_r} 2(p_{ik} - \delta_{ik} p_{jj}) dv^i.$$

(E, P_1, P_2, P_3) is the so called ADM energy momentum 4-vector.

The positive mass theorem of Schoen and Yau^{21,22} (see also Witten³²) states that the total mass of such an isolated system is always positive. Suppose the dominant energy condition holds along an asymptotically flat Ω , then (E, P_1, P_2, P_3) is a

future-directed non-spacelike vector, i.e.

$$E \geq 0, -E^2 + P_1^2 + P_2^2 + P_3^2 \leq 0.$$

In particular, the ADM mass $\sqrt{E^2 - P_1^2 - P_2^2 - P_3^2}$ is non-negative and $= 0$ if and only if the spacetime is flat along $\bar{\Omega}$.

There is also the Bondi-Sachs energy-momentum² for a asymptotically null hypersurface which measures energy after radiation. Positive energy theorem at null infinity also holds,^{12,24} and thus the physical system cannot radiate away more energy than it has initially.

4. Quasilocal energy/mass and expectations

We formulate the question of quasilocal energy and mass:

Question 4.1. Suppose Ω is a bounded spacelike region, what is the total energy intercepted by Ω as seen by an observer? What is the total mass contained in Ω ? The answer to these questions should depend only on $\Sigma = \partial\Omega$ by conservation law.

In comparison to the ADM or Bondi total mass for an isolated system where gravitation is weak at boundary (infinity), the notion of quasilocal mass corresponds to a non-isolated system where gravitation could be strong. What properties qualify for a valid definition? Here are three that we think are most natural:

- (1) Asymptotics: The limit should recover the ADM mass in the asymptotically flat case and the Bondi mass in the asymptotically null case. It should also recover the energy-momentum tensor in non-vacuum and the Bel-Robinson tensor in vacuum for small sphere limits.
- (2) Positivity: The mass should be positive under local energy condition for a large class of surfaces.
- (3) Rigidity: The quasilocal mass should vanish for surfaces in $\mathbb{R}^{3,1}$.

5. Hamilton-Jacobi approach

There have been various approaches in attempt to define quasilocal mass (see²⁶ and the reference therein). We focus on the canonical Hamilton-Jacobi analysis approach which seems most relevant to Einstein's equation. Quasilocal Hamilton-Jacobi analysis of Einstein-Hilbert action has been studied by Brown-York,^{5,6} Hawking-Horowitz,¹¹ and Kijowski.¹³ Applying the analysis to the time history of a spatially bounded region in spacetime yields the Hamiltonian which is a 2-surface integral at terminal time that depends on a pair of vector fields (t^μ, u^μ) along Σ . t^μ is a future timelike unit vector field and u^μ a future timelike unit normal vector. u^μ should be considered as the future unit normal of a spacelike hypersurface Ω bounded by Σ . We decompose

$$t^\mu = Nu^\mu + N^\mu.$$

The surface Hamiltonian in¹¹ is

$$\mathcal{H}(t^\mu, u^\mu) = -\frac{1}{8\pi} \int_{\Sigma} Nk - N^\mu v^\nu (p_{\mu\nu} - p^\lambda_\lambda g_{\mu\nu}) \quad (3)$$

where k is the mean curvature of Σ as boundary of Ω , $p_{\mu\nu}$ is the second fundamental form of Ω in spacetime, and v^ν is the outward unit spacelike normal along Σ that is orthogonal to u^ν .

The energy is defined to be the difference between the physical surface Hamiltonian and the reference surface Hamiltonian. Reference surface Hamiltonian in principle should come from data associated with isometric embedding of the time history of the boundary into a reference spacetime. But this is in general an over-determined problem.

Isometric embedding of Σ into \mathbb{R}^3 has been used to define Brown-York mass and Liu-Yau mass (see also Kijowski,¹³ Booth-Mann,³ Epp,⁹ etc.) with $u^\mu = t^\mu$ (thus $N = 1$ and $N^\mu = 0$) to be specified. There is a unique isometric embedding into \mathbb{R}^3 for any metric with positive Gauss curvature, see Nirenberg¹⁹ and Pogorelov.²⁰

The Brown-York mass is defined to be $\frac{1}{8\pi}(\int_{\Sigma} k_0 - \int_{\Sigma} k)$ where k is the mean curvature of Σ with respect to a spacelike region Ω , and k_0 is the mean curvature of the image of the isometric embedding of Σ into \mathbb{R}^3 . The Liu-Yau mass is $\frac{1}{8\pi}(\int_{\Sigma} k_0 - \int_{\Sigma} |H|)$ where H is the mean curvature vector of Σ in spacetime. Note that the Liu-Yau mass is gauge independent.

The Brown-York mass and the Liu-Yau mass have the important positivity property by the work of Shi-Tam²⁵ and Liu-Yau,^{14,15} respectively. However, there exist surfaces in $\mathbb{R}^{3,1}$ with strictly positive Brown-York mass and Liu-Yau mass.¹⁷

6. New definition of quasilocal energy

For an isometric embedding $X : \Sigma \rightarrow \mathbb{R}^{3,1}$ and $T_0 \in \mathbb{R}^{3,1}$ a constant future timelike unit vector, we define the quasilocal energy to be

$$E(\Sigma, X, T_0) = \mathcal{H}(t^\mu, u^\mu) - \mathcal{H}(t_0^\mu, u_0^\mu)$$

where $t_0^\mu = T_0$.

We shall call $\Sigma \subset M$ the physical surface and the image of X in $\mathbb{R}^{3,1}$ the reference surface.

In the following, we discuss our prescription for u_0^μ , t^μ , and u^μ in.^{27,28} Consider the reference surface $\Sigma \subset \mathbb{R}^{3,1}$ and t_0^μ a constant future timelike unit vector. We take u_0^μ to be the unit normal future timelike unit vector field in the direction of the normal part of t_0^μ , i.e. $t_0^\mu = Nu_0^\mu + N^\mu$ where N^μ is tangent to Σ .

This defines the reference Hamiltonian $\mathcal{H}(t_0^\mu, u_0^\mu)$ which is shown to be equal to

$$-\frac{1}{8\pi} \int_{\hat{\Sigma}} \hat{k}$$

where $\hat{\Sigma}$ is the projection of Σ onto the orthogonal complement of $t_0^\mu = T_0$.

We proved a unique isometric embedding theorem²⁸ into $\mathbb{R}^{3,1}$ with convex shadows, i.e. $\hat{\Sigma}$ is a convex surface in the orthogonal complement \mathbb{R}^3 . To find the corresponding gauge (t^μ, u^μ) on the physical surface, we assume the mean curvature vector of Σ in spacetime is spacelike. For a reference isometric embedding $X : \Sigma \rightarrow \mathbb{R}^{3,1}$ and a t_0^μ , we claim there exists a unique future timelike unit vector t^μ along the physical surface $\Sigma \subset M$ such that

“The expansion of Σ along t_0^μ in $\mathbb{R}^{3,1}$ is the same as the expansion of Σ along t^μ in M ”.

Now define u^μ by $t^\mu = Nu^\mu + N^\mu$ along the physical surface $\Sigma \subset M$ for the same N and N^μ . Thus t^μ and t_0^μ have the same lapse functions and shift vectors along the physical surface $\Sigma \subset M$ and the reference surface $\Sigma \subset \mathbb{R}^{3,1}$, respectively. Use this (t^μ, u^μ) on $\Sigma \subset M$ to compute the physical Hamiltonian $\mathcal{H}(t^\mu, u^\mu)$ and this defines our quasilocal energy $E(\Sigma, X, T_0)$.

7. The expression and properties

Let Σ be a spacelike 2-surface in spacetime which bounds a spacelike hypersurface Ω with a future unit timelike normal vector field u^μ . Denote by v^μ the unit spacelike outward normal of $\Sigma = \partial\Omega$ with respect to Ω . The mean curvature vector of Σ is

$$H = -kv^\mu + pu^\mu$$

where k is the mean curvature of Σ in Ω with respect to v^μ and p is the trace of the restriction of p_{ij} to Σ . The definition of H is indeed independent of Ω and the choice of u^μ and v^μ . Let J be the reflection of H along the future inward null direction in the normal bundle, i.e. $J = ku^\mu - pv^\mu$. H is inward spacelike if and only if J is future timelike.

It turns out $E(\Sigma, X, T_0)$ can be expressed in term of the mean curvature vector field H of Σ in M and $\tau = -\langle X, T_0 \rangle_{\mathbb{R}^{3,1}}$.

Suppose H is spacelike, we can use the frame H and J to define a connection one-form for the normal bundle of Σ by $\langle \nabla_{(\cdot)}^M \frac{J}{|H|}, \frac{H}{|H|} \rangle$. We recall the following fact that *“the mean curvature vector of the isometric embedding $X : \Sigma \rightarrow \mathbb{R}^{3,1}$ is $H_0 = \Delta X$ ”*. Here Δ is the Laplace operator for functions on Σ with respect to the induced metric. For a function defined on Σ such as τ , we also use $\nabla\tau$ to denote its gradient vector that is tangent to Σ . The quasilocal energy $E(\Sigma, X, T_0)$ with respect to (X, T_0) is

$$\begin{aligned} & \frac{1}{8\pi} \int_{\hat{\Sigma}} \hat{k} - \frac{1}{8\pi} \int_{\Sigma} [\sqrt{|H|^2(1 + |\nabla\tau|^2) + (\Delta\tau)^2} \\ & - \Delta\tau \sinh^{-1} \frac{\Delta\tau}{\sqrt{1 + |\nabla\tau|^2}|H|} - \langle \nabla_{\nabla\tau}^M \frac{J}{|H|}, \frac{H}{|H|} \rangle] \end{aligned}$$

where

$$\begin{aligned} & \int_{\hat{\Sigma}} \hat{k} = \int_{\Sigma} [\sqrt{|H_0|^2(1 + |\nabla\tau|^2) + (\Delta\tau)^2} \\ & - \Delta\tau \sinh^{-1} \frac{\Delta\tau}{\sqrt{1 + |\nabla\tau|^2}|H_0|} - \langle \nabla_{\nabla\tau}^{\mathbb{R}^{3,1}} \frac{J_0}{|H_0|}, \frac{H_0}{|H_0|} \rangle]. \end{aligned}$$

Quasilocal mass is defined to be the infimum of quasilocal energy $E(\Sigma, X, T_0)$ among all “admissible observers” (X, T_0) (see²⁷ for the definition):

$$m(\Sigma) = \inf E(\Sigma, X, T_0).$$

In,^{7,27–29} we prove:

- (1) Positivity: $m(\Sigma) \geq 0$ under dominant energy condition on spacetime and convexity assumptions on Σ .
- (2) Rigidity: $m(\Sigma) = 0$ if Σ is in $\mathbb{R}^{3,1}$.
- (3) Quasilocal mass approaches the ADM mass and Bondi mass at spatial and null infinity, respectively.

This is the only known definition of quasilocal mass that satisfies all these properties.

In fact, the quasilocal energy $E(S_r, X_r, T_0)$ gets linearized and acquires the Lorentzian symmetry at infinity.

$$\lim_{r \rightarrow \infty} E(S_r, X_r, T_0) = T_0^\mu P_\mu$$

where $P_\mu = (P_0, P_1, P_2, P_3)$ is the ADM / Bondi-Sachs energy-momentum 4-vector, at spatial/null infinity.

In general, suppose Σ_r is a family of surface in spacetime and a family of isometric embedding X_r of Σ_r in $\mathbb{R}^{3,1}$ is given. As long as $\frac{|H_0|}{|H|} \rightarrow 1$ as $r \rightarrow \infty$, the limit of the quasilocal energy $E(\Sigma_r, X_r, T_0)$ is the same as the limit of

$$\frac{1}{8\pi} \int_{\Sigma_r} -\langle T_0, \frac{J_0}{|H_0|} \rangle (|H_0| - |H|) - \langle \nabla_{\nabla^\tau}^{\mathbb{R}^{3,1}} \frac{J_0}{|H_0|}, \frac{H_0}{|H_0|} \rangle + \langle \nabla_{\nabla^\tau}^N \frac{J}{|H|}, \frac{H}{|H|} \rangle.$$

As $\tau = -\langle X, T_0 \rangle$, the expression is already linear in T_0 .

8. Surface Hamiltonian and Minkowski inequality

In this section, we discuss the surface Hamiltonian in Minkowski space and the connection to an inequality proposed by Gibbons.

We recall the following identity in^{28,30} regarding the surface Hamiltonian (3):

Proposition 8.1. *For a closed spacelike 2-surface Σ in the Minkowski space which bounds a spacelike hypersurface and a constant future timelike unit vector field T_0 , there exists a unique orthogonal normal gauge $\{\check{e}_3, \check{e}_4\}$ along Σ such that \check{e}_3 is a outward spacelike unit normal and \check{e}_4 is a future timelike unit normal and they satisfy*

$$-\frac{1}{8\pi} \int_{\Sigma} \langle J, T_0 \rangle_{\mathbb{R}^{3,1}} + \langle \nabla_{\check{e}_3}^{\mathbb{R}^{3,1}} \check{e}_4, T_0^\top \rangle_{\mathbb{R}^{3,1}} = \frac{1}{8\pi} \int_{\hat{\Sigma}} \hat{k} \quad (4)$$

where $\hat{\Sigma}$ is the projection of Σ onto the orthogonal complement of T_0 and \hat{k} is the mean curvature of $\hat{\Sigma}$.

Proof. Proposition 3.1 of²⁸ (see also¹⁰). \square

In fact, denote by τ the restriction of time function defined by T_0 to Σ and by $\nabla\tau$ the gradient vector field of τ on Σ with respect to the induced metric, we have

$$T_0 = \sqrt{1 + |\nabla\tau|^2} \check{e}_4 - \nabla\tau.$$

The lapse and shift of T_0 are given by $\sqrt{1 + |\nabla\tau|^2}$ and $T_0^\top = -\nabla\tau$, respectively.

The classical Minkowski inequality for surfaces in \mathbb{R}^3 states that for a closed convex surface $\hat{\Sigma}$ in \mathbb{R}^3 ,

$$\int_{\hat{\Sigma}} \hat{k} d\mu \geq \sqrt{16\pi |\hat{\Sigma}|},$$

where \hat{k} is the mean curvature and $|\hat{\Sigma}|$ is the area of $\hat{\Sigma}$.

Applying the Minkowski inequality and recalling that the area of $\hat{\Sigma}$ is always greater than or equal to the area of Σ , we obtained the following inequality between the surface Hamiltonian and the area.

$$-\frac{1}{8\pi} \int_{\Sigma} \langle J, T_0 \rangle_{\mathbb{R}^{3,1}} + \langle \nabla_{\check{e}_3}^{\mathbb{R}^{3,1}} \check{e}_4, T_0^\top \rangle_{\mathbb{R}^{3,1}} \geq \sqrt{\frac{|\Sigma|}{4\pi}}. \quad (5)$$

In equation (6.16) of,¹⁰ the author claimed that the following inequality holds and called it the black hole isoperimetric inequality.

$$-\frac{1}{8\pi} \int_{\Sigma} \langle J, T_0 \rangle_{\mathbb{R}^{3,1}} \geq \sqrt{\frac{|\Sigma|}{4\pi}}. \quad (6)$$

However, the derivation in¹⁰ is not correct and the validity of this inequality remains open, see also section 7.1 of.¹⁶

Recently, a sharp Minkowski inequality in the hyperbolic 3-space was proved in:⁴

$$\int_{\Sigma} f h d\mu - 6 \int_{\Omega} f d\text{vol} \geq \sqrt{16\pi |\Sigma|} \quad (7)$$

for any mean convex, star shaped region $\Omega \subset \mathbb{H}^3$ and $\Sigma = \partial\Omega$. Here $f = \cosh r$ where r is the geodesic distance function with respect to a point $o \in \mathbb{H}^3$ and h is the mean curvature of Σ with respect to outward unit normal of Ω .

In the following, we show that (7) is equivalent to (6) when the surface $\Sigma \subset \mathbb{R}^{3,1}$ lies in the hyperbolic space $\mathbb{H}^3 \subset \mathbb{R}^{3,1}$.

Theorem 8.1. *Suppose Σ is a closed embedded spacelike 2-surface in the Minkowski space and $T_0 \in \mathbb{H}^3 \subset \mathbb{R}^{3,1}$ is a future unit timelike constant vector. Suppose Σ lies in \mathbb{H}^3 and is mean convex and star-shaped with respect to T_0 , then inequality (6) holds, i.e.*

$$-\frac{1}{8\pi} \int_{\Sigma} \langle J, T_0 \rangle_{\mathbb{R}^{3,1}} \geq \sqrt{\frac{|\Sigma|}{4\pi}}.$$

Proof. By Lorentz transformation, we may assume T_0 is $(1, 0, 0, 0)$. Now choose r to be the geodesic distance function on \mathbb{H}^3 with $(1, 0, 0, 0)$ as the origin. We embed \mathbb{H}^3 into $\mathbb{R}^{3,1}$ as the upper branch of the hyperbola $\{(t, x, y, z) \mid t > 0, -t^2 + x^2 + y^2 + z^2 = -1\}$. Suppose the embedding is given by the position four-vector $X = (t, x, y, z)$

The mean curvature vector of Σ in $\mathbb{R}^{3,1}$ is $-h\nu + 2e_4$ where h is the mean curvature of Σ in \mathbb{H}^3 , ν is the outward unit normal of Σ in \mathbb{H}^3 , and e_4 is the future unit timelike normal of \mathbb{H}^3 in $\mathbb{R}^{3,1}$. Note that e_4 is the same as the position vector of the embedding $X : \mathbb{H}^3 \rightarrow \mathbb{R}^{3,1}$.

Consider the normal vector field

$$J = he_4 - 2\nu$$

obtained by reflecting H along the future inward null direction of the normal bundle. We check that

$$-\int_{\Sigma} \langle J, T_0 \rangle_{\mathbb{R}^{3,1}} d\mu = \int_{\Sigma} (f h - 2 \frac{\partial f}{\partial \nu}) d\mu = \int_{\Sigma} f h d\mu - 6 \int_{\Omega} f d\text{vol}$$

because $f = -\langle X, T_0 \rangle$ and $\bar{\Delta}f = 3f$ where $\bar{\Delta}$ is the Laplace operator on \mathbb{H}^3 . \square

If we assume the surface has spacelike inward mean curvature vector, the integral $-\int_{\Sigma} \langle J, T_0 \rangle_{\mathbb{R}^{3,1}} d\mu$ is positive for any T_0 . From this, we can formulate a general question for spacelike surfaces in $\mathbb{R}^{3,1}$.

Question 8.1. Suppose Σ is closed spacelike 2-surface that bounds a spacelike hypersurface in $\mathbb{R}^{3,1}$. Under what condition does inequality (6) hold?

Acknowledgements

The author would like to thank Simon Brendle, PoNing Chen, Gary Gibbons, and Shing-Tung Yau for helpful discussions. The author's research was supported by the National Science Foundation under grant DMS-1105483.

References

1. R. Arnowitt, S. Deser, and C. W. Misner, *The dynamics of general relativity*, 1962 Gravitation: An introduction to current research pp. 227-265 Wiley, New York. arXiv:gr-qc/0405109
2. H. Bondi, M. G. van der Burg and A. W. K. Metzner, *Gravitational waves in general relativity. VII. Waves from axi-symmetric isolated systems*, Proc. Roy. Soc. Ser. A, **269** (1962) 21-52.
3. I. S. Booth and R. B. Mann, *Moving observers, nonorthogonal boundaries, and quasilocal energy*, Phys. Rev. D, **59**, 064021 (1999).
4. S. Brendle, P.-K. Hung, and M.-T. Wang, *A Minkowski type inequality for hypersurfaces in the Anti-deSitter-Schwarzschild manifold*, arXiv:1209.0669
5. J. D. Brown and J. W. York, *Quasilocal energy in general relativity*, Mathematical aspects of classical field theory (Seattle, WA, 1991), 129-142, Contemp. Math. **132**, Amer. Math. Soc., Providence, RI, 1992.

6. J. D. Brown and J. W. York, *Quasilocal energy and conserved charges derived from the gravitational action*, Phys. Rev. D (3) **47** (1993), no. 4, 1407–1419.
7. P. Chen, M.-T. Wang, and S.-T. Yau, *Evaluating quasilocal energy and solving optimal embedding equation at null infinity*, Comm. Math. Phys. **308** (2011), no.3, 845–863.
8. D. Christodoulou, *Mathematical problems of general relativity. I*, Zürich Lectures in Advanced Mathematics. European Mathematical Society (EMS), Zürich, 2008. x+147 pp.
9. R. J. Epp, *Angular momentum and an invariant quasilocal energy in general relativity*, Phys. Rev. D **62** (2000), no. 12, 124108.
10. G. W. Gibbons, *Collapsing shells and the isoperimetric inequality for black holes*, Classical Quantum Gravity **14** (1997), no. 10, 2905–2915.
11. S. W. Hawking and G. T. Horowitz, *The gravitational Hamiltonian, action, entropy and surface terms*, Classical Quantum Gravity **13** (1996), no. 6, 1487–1498.
12. G. T. Horowitz and M. J. Perry, *Gravitational energy cannot become negative*, Phys. Rev. Lett. **48** (1982), no. 6, 371–374.
13. J. Kijowski, *A simple derivation of canonical structure and quasi-local Hamiltonians in general relativity*, Gen. Relativity Gravitation **29** (1997), no. 3, 307–343.
14. C.-C. M. Liu and S.-T. Yau, *Positivity of quasilocal mass*, Phys. Rev. Lett. **90** (2003), no. 23, 231102.
15. C.-C. M. Liu and S.-T. Yau, *Positivity of quasilocal mass II*, J. Amer. Math. Soc. **19** (2006), no. 1, 181–204.
16. M. Mars, *Present status of the Penrose inequality*, Classical Quantum Gravity **26** (2009), no. 19, 193001, 59 pp.
17. N. Ó Murchadha, L. B. Szabados, and K. P. Tod, *Comment on “Positivity of quasilocal mass”*, Phys. Rev. Lett. **92** (2004), 259001.
18. R. Penrose, *Some unsolved problems in classical general relativity*, Seminar on Differential Geometry, pp. 631–668, Ann. of Math. Stud., 102, Princeton Univ. Press, Princeton, N.J., 1982.
19. L. Nirenberg, *The Weyl and Minkowski problems in differential geometry in the large*, Comm. Pure Appl. Math. **6**, (1953). 337–394.
20. A. V. Pogorelov, *Regularity of a convex surface with given Gaussian curvature*, (Russian) Mat. Sbornik N.S. **31**(73), (1952). 88–103.
21. R. Schoen and S.-T. Yau, *Positivity of the total mass of a general space-time*, Phys. Rev. Lett. **43** (1979), no. 20, 1457–1459.
22. R. Schoen and S.-T. Yau, *On the proof of the positive mass conjecture in general relativity*, Comm. Math. Phys. **65** (1979), no. 1, 45–76.
23. R. Schoen and S.-T. Yau, *Proof of the positive mass theorem II*, Comm. Math. Phys. **79** (1981), no. 2, 231–260.
24. R. Schoen and S.-T. Yau, *Proof that the Bondi mass is positive*, Phys. Rev. Lett. **48** (1982), no. 6, 369–371.
25. Y. Shi and L.-F. Tam, *Positive mass theorem and the boundary behavior of compact manifolds with nonnegative scalar curvature*, J. Differential Geom. **62** (2002), no. 1, 79–125.
26. L. B. Szabados, *Quasi-Local Energy-Momentum and Angular Momentum in GR: A Review Article*, Living Rev. Relativity, **12** (2009) No 4, URL: relativity.livingreviews.org/Articles/lrr-2009-4
27. M.-T. Wang and S.-T. Yau, *Quasilocal mass in general relativity*, Phys. Rev. Lett. **102** (2009), no. 2, no. 021101.
28. M.-T. Wang and S.-T. Yau, *Isometric embeddings into the Minkowski space and new quasi-local mass*, Comm. Math. Phys. **288** (2009), no. 3, 919–942.

29. M.-T. Wang and S.-T. Yau, *Limit of quasilocal mass at spatial infinity*, Comm. Math. Phys. **296** (2010), no.1, 271–283.
30. M.-T. Wang, *Quasilocal mass from a mathematical perspective*, Fifth International Congress of Chinese Mathematicians. Part 1, 2, 355364, AMS/IP Stud. Adv. Math., 51, pt. 1, 2, Amer. Math. Soc., Providence, RI, 2012.
31. M.-T. Wang, *Gravitational energy seen by quasilocal observers*, Classical Quantum Gravity **28** (2011), no. 11, 114011, 9 pp.
32. E. Witten, *A new proof of the positive energy theorem*, Comm. Math. Phys. **80** (1981), no. 3, 381–402.